

Cosmic Dawn Intensity Mapper CDIM


Tzu-Ching Chang
(Jet Propulsion Laboratory,
California Institute of Technology)

for the


CDIM Science Team

and

CDIM Design Team



Cosmic Dawn Intensity Mapper (CDIM)



Asantha Cooray, UC Irvine (PI)

Tzu-Ching Chang, JPL (Study Scientist)

Stephen Unwin, JPL (Study Manager)

Science Team

Renyue Cen, Princeton

Ranga Ram Chary, IPAC

Olivier Doré, JPL

Xiaohui Fan, Arizona

Giovanni Fazio, CfA

Chang Feng, U Illinois

Steven Finkelstein, U T Austin

Hai Fu, U Iowa

Caroline Heneka, SNS

Hooshang Nayyeri, UC Irvine

Jason Rhodes, JPL

Mario Santos, U Western Cape

Raphael Sadoun, Osaka U

Marta Silva, U Oslo

Hy Trac, CMU

Heidi Wu, OSU

Michael Zemcov, RIT

Zheng Zheng, U Utah

Design Team

Patrick Morrissey, JPL
(Instrument)

Andrew Coffey, JPL (System
engineering)

Nasrat Raouf, JPL (LVF filters)

Talso Chui, JPL (Thermal)

James McGuire, JPL (Optics)

Sarah Lipsky, Ball (Bus)


CDIM in a (Falcon-9) Nutshell

NASA Probe class mission concept for the 2020 Decadal review.


- 0.75 μm – 7.5 μm spectro-imaging in 860 narrow bands at $R=300$
- 0.8 m effective aperture
- 7.7 sq. degree focal plane
- Diffraction limited at 7.5 μm , 2" PSF
- Three-tiered survey in 4 years
- Costed at JPL under I\$B (incl. 30% margin)

Cosmic Dawn and Reionization Sciences

- First Galaxies: tracing $\text{H}\alpha$ to $z=10$
- First Blackholes: finding AGNs at $z=8$
- Reionization Tomography: $\text{Ly}\alpha$, $\text{H}\alpha$ intensity mapping, and cross-correlation with 21 cm EoR maps




Cosmic Dawn and Reionization



NASA


Cosmic Dawn and Reionization



Galaxies




AGN



IGM


CDIM Holistic View of Cosmic Dawn and Reionization



Galaxies:
Measuring H α up
to $z \sim 10$





AGNs:
Finding
blackholes up to
 $z \sim 8$



IGM Tomography:
reionization
topology and
history

CDIM Holistic View of Cosmic Dawn and Reionization




Galaxies:
Measuring H α up to $z \sim 10$
Wavelength 0.75-7.5 microns
 $R \geq 300$ to separate [NII] &
Line flux sensitivity

AGNs:
Finding
blackholes up to $z \sim 8$
Large survey area &
point source sensitivity

IGM Tomography:
Reionization topology and
history
Large field of view &
Surface brightness sensitivity

CDIM Design

5.6m



Three-mirror all-reflective design with 0.8m clear aperture


Linear Variable Filters (LVF) at $R=300$, 4x6 H2RG detectors

V-groove radiators, passive cooling at $T < 35\text{K}$ in L2 halo orbit

Ka-band HGA. Data rate ~ 400 Gbit/day, 1hr/day downlink


Design team & Team-X

H2RG detectors, Linear Variable Filters (LVF) and the FoV



- HyViSi and H2RGs in 4x6 array across 0.75-7.5 microns; at R=300 resulting in 840 wavelength channels.
- $2.3 \times 3.4 = 7.7 \text{ deg}^2$ FoV

Spectroscopy with Linear Variable Filters



Construct 3D spectral-imaging data cube at 1'' pixels in 840 wavelength bands between 0.75 μm – 7.5 μm

- Build up wavelength coverage by stepping spatially, each step is 15''
- Each exposure/step is 250s (ZL-limited), 5 days to build up a FoV-worth of full data cube

SPHEREx


CDIM Three-tiered Survey



- Detecting faint, high-redshift galaxies
 - Deep survey; 15 deg² to overlap with WFIRST/Euclid deep fields




- Detecting bright, rare AGNs
 - Wide survey; 300 deg² to catch z=8 AGNs.
 - At either SEP or NEP, visible all-year round from L2, surrounding the Deep survey field



- Reionization tomography in synergy with 21cm intensity maps
 - Medium survey; 30 deg² to match a SKA 21cm EoR deep field likely overlapping with the ECDF-S and HERA
- Four years of survey




CDIM Point-Source Sensitivity







CDIM Line-flux Sensitivity



CDIM R=300
840 bands



CDIM Surface Brightness Sensitivity



CDIM R=300
840 bands

CDIM R=3
broad band



CDIM Traces H α up to $z \sim 10$

Preliminary!

deep
Medium
Wide

deep
Medium
Wide

deep
Medium
Wide




Heidi Hao-Yi Wu

- Optimistic and pessimistic H α luminosity functions based on Flexible Stellar Population Synthesis code (Conroy 2010), Millennium simulation, UV galaxy measurements and semi-analytic models
- Optimistic scenario leads to H α detections up to $z=10$

deep
Medium
Wide


Measuring Stellar Mass and Metal production



Steve Finkelstein


- Infer SFR from rest-frame UV and metallicity from $[NII]/H\alpha$.
- Trace stellar mass and metal build up across redshift

Finding AGNs out to $z \sim 8$



- Two quasar luminosity functions: based on Willott+15 and Kulkarni+18 and Wang+18, which has a steeper faint-end slope
- Possibly finding >10 quasars at $z=7$ in CDIM Wide Survey

Estimating Blackhole Masses




Onoue+18

Renyue Cen


- CDIM will find ~ 100 quasars at $z=6$. Detection limit corresponds to a blackhole mass estimate of $10^7 M_{\text{sun}}$, assuming Eddington accretion.

Detecting Ly α Emitting Galaxies

LAE luminosity function



LAE number count




Zheng Zheng

- Detect a few thousand LAEs with $L_{\text{Ly}\alpha} > 10^{43}$ erg/s
- Determine the bright end of Ly α luminosity function at $5.2 < z < 7$
- H α counter part detectable out to $z \sim 9$, can stack on H α for fainter LAEs.
- CDIM spectral resolution ~ 6 Mpc/h at $z=6$ v.s. Silverrush narrowband at ~ 30 Mpc/h
- Constrain the evolution of ionization state of IGM

Fainter Ly α Emitters


Stacking Ly α images around
H α selected galaxies with CDIM



Raphael Sadoun




- Stacked LAE images using (83) CDIM detected H α galaxies at $z=5.7$ within a 50^3 Mpc/h simulation volume
- Ly α images predicted by Ly α ray-tracing model + reionization simulations

21cm, H α , Ly α Reionization Tomography






21cm, H α , Ly α auto and cross-power spectra



- H α (and Ly α) trace ionizing sources. Ly α scatter in the ionized IGM. Cross-correlation can probe the scales of Ly α photon scattering in the IGM during EoR.
- 21cm x H α and 21cm x Ly α are anti-correlated on typical scale of ionized regions. Can be used to trace bubble growth as a function of redshift.



- Cross-correlations are more robust against potential systematics. Verify 21cm cosmological signals.



Measuring Reionization History with CDIM



Marta Silva

- Ionization fraction can be inferred from multiple measurements: evolution of LAEs, H α and Ly α intensity evolution, and amplitudes of (21 cm \times Ly α) cross-power spectra.

Summary

- CDIM probe design developed:
 - 0.75 μm – 7.5 μm spectro-imaging at $R=300$
 - 0.8 m effective aperture, 1" pixel, 2" PSF
 - 7.7 sq. degree focal plane
 - Three-tiered survey in 4 years (15, 30, 300 sq. degree)
 - Costed with Class-B margin
- Designed to probe Cosmic Dawn and Reionization in novel and powerful ways:
 - First Galaxies: tracing $\text{H}\alpha$ to $z=10$, studying stellar mass and metal build up
 - First Blackholes: finding AGNs at $z=8$, constraining blackhole mass growth
 - Reionization Tomography: $\text{Ly}\alpha$, $\text{H}\alpha$ Intensity Mapping, cross-correlation with 21cm, to measure bubble growth and reionization history
- CDIM will deliver a unique data set and offer a holistic view of Cosmic Dawn and Reionization